



## Weighted Poincaré Inequalities and Applications in Domain Decomposition

Clemens Pechstein

Institute of Computational Mathematics, Johannes Kepler University  
Altenberger Str. 69, 4040 Linz, Austria

Robert Scheichl

Department of Mathematical Sciences, University of Bath,  
Claverton Down, Bath BA2 7AY, UK

NuMa-Report No. 2009-12

November 2009

## Technical Reports before 1998:

### 1995

- 95-1 Hedwig Brandstetter  
*Was ist neu in Fortran 90?* March 1995
- 95-2 G. Haase, B. Heise, M. Kuhn, U. Langer  
*Adaptive Domain Decomposition Methods for Finite and Boundary Element Equations.* August 1995
- 95-3 Joachim Schöberl  
*An Automatic Mesh Generator Using Geometric Rules for Two and Three Space Dimensions.* August 1995

### 1996

- 96-1 Ferdinand Kickingger  
*Automatic Mesh Generation for 3D Objects.* February 1996
- 96-2 Mario Goppold, Gundolf Haase, Bodo Heise und Michael Kuhn  
*Preprocessing in BE/FE Domain Decomposition Methods.* February 1996
- 96-3 Bodo Heise  
*A Mixed Variational Formulation for 3D Magnetostatics and its Finite Element Discretisation.* February 1996
- 96-4 Bodo Heise und Michael Jung  
*Robust Parallel Newton-Multilevel Methods.* February 1996
- 96-5 Ferdinand Kickingger  
*Algebraic Multigrid for Discrete Elliptic Second Order Problems.* February 1996
- 96-6 Bodo Heise  
*A Mixed Variational Formulation for 3D Magnetostatics and its Finite Element Discretisation.* May 1996
- 96-7 Michael Kuhn  
*Benchmarking for Boundary Element Methods.* June 1996

### 1997

- 97-1 Bodo Heise, Michael Kuhn and Ulrich Langer  
*A Mixed Variational Formulation for 3D Magnetostatics in the Space  $H(\text{rot}) \cap H(\text{div})$*  February 1997
- 97-2 Joachim Schöberl  
*Robust Multigrid Preconditioning for Parameter Dependent Problems I: The Stokes-type Case.* June 1997
- 97-3 Ferdinand Kickingger, Sergei V. Nepomnyaschikh, Ralf Pfau, Joachim Schöberl  
*Numerical Estimates of Inequalities in  $H^{\frac{1}{2}}$ .* August 1997
- 97-4 Joachim Schöberl  
*Programmbeschreibung NAOMI 2D und Algebraic Multigrid.* September 1997

From 1998 to 2008 technical reports were published by SFB013. Please see

<http://www.sfb013.uni-linz.ac.at/index.php?id=reports>

From 2004 on reports were also published by RICAM. Please see

<http://www.ricam.oeaw.ac.at/publications/list/>

For a complete list of NuMa reports see

<http://www.numa.uni-linz.ac.at/Publications/List/>

# Weighted Poincaré Inequalities and Applications in Domain Decomposition

Clemens Pechstein and Robert Scheichl

**Abstract** Poincaré type inequalities play a central role in the analysis of domain decomposition and multigrid methods for second-order elliptic problems. However, when the coefficient varies within a subdomain or within a coarse grid element, then standard condition number bounds for these methods may be overly pessimistic. In this short note we present new weighted Poincaré type inequalities for a class of piecewise constant coefficients that lead to sharper bounds independent of any possible large contrasts in the coefficients.

## 1 Introduction

Poincaré type inequalities play a central role in the analysis of domain decomposition (DD) methods for finite element discretisations of elliptic PDEs of the type

$$-\nabla \cdot (\alpha \nabla u) = f. \quad (1)$$

In many applications the coefficient  $\alpha$  in (1) is discontinuous and varies over several orders of magnitude throughout the domain in a possibly very complicated way. Standard analyses of DD methods for (1) that use classical Poincaré type inequalities will often lead to pessimistic bounds. Two examples are the popular two-level overlapping Schwarz and FETI. If the subdomain partition can be chosen such that  $\alpha$  is constant (or almost constant) on each subdomain as well as in each element of

---

Clemens Pechstein  
Institute of Computational Mathematics, Johannes Kepler University Linz, Altenberger Str. 69,  
4040 Linz, Austria, e-mail: Clemens.Pechstein@jku.at,  
supported by the Austrian Sciences Fund (FWF) under grant P19255

Robert Scheichl  
Department of Mathematical Sciences, University of Bath, Claverton Down, Bath BA2 7AY, UK,  
e-mail: R.Scheichl@bath.ac.uk

the coarse mesh (for two-level methods), then it is possible to prove bounds that are independent of the coefficient variation (cf. [3, 8, 16]). However, if this is not possible and the coefficient varies strongly within a subdomain, then the classical bounds depend on the local variation of the coefficient, which may be overly pessimistic in many cases. To obtain sharper bounds in some of these cases, it is possible to refine the standard analyses and use Poincaré inequalities on annulus type boundary layers of each subdomain [6, 10, 12, 14, 9], or weighted Poincaré type inequalities [5, 11, 15]. See also [2, 4, 7, 13, 18] for related work.

In this short note we want to collect and expand on the results in [5, 11] and present a new class of weighted Poincaré-type inequalities for a rather general class of piecewise constant coefficients. Due to space restrictions we have to refer the interested reader to [6, 10, 11, 15], to see where exactly these new inequalities can be used in the analysis of FETI and two-level Schwarz methods. Note that the inequalities are much more widely applicable, e.g. in the analysis of multigrid methods for (1). In particular they can be used to improve the results in [17], which rely on estimates on the weighted  $L_2$ -projection in [1] and therefore require the coefficient to be resolved by the coarse grids.

## 2 Weighted Poincaré Inequalities and Discrete Analogues

Let  $D$  be a bounded domain in  $\mathbb{R}^d$ ,  $d = 2, 3$ . For simplicity we only consider piecewise constant coefficient functions  $\alpha$  with respect to a non-overlapping partitioning of  $D$  into open, connected Lipschitz polygons (polyhedra)  $\{Y_\ell : \ell = 1, \dots, n\}$ , i.e.  $\alpha|_{Y_\ell} \equiv \alpha_\ell$  for some constants  $\alpha_\ell$ . The results generalise in a straightforward way to more general coefficients that vary mildly within each of the regions  $Y_\ell$ .


**Definition 1.** The region  $P_{\ell_1, \ell_s} := (\bar{Y}_{\ell_1} \cup \bar{Y}_{\ell_2} \cup \dots \cup \bar{Y}_{\ell_s})^\circ$  is called a *type- $m$  quasi-monotone path* from  $Y_{\ell_1}$  to  $Y_{\ell_s}$ , if the following two conditions are satisfied:

- (i)  $\bar{Y}_{\ell_i}$  and  $\bar{Y}_{\ell_{i+1}}$  share a common  $m$ -dimensional manifold  $X_i$ , for  $i = 1, \dots, s-1$ ,
- (ii)  $\alpha_{\ell_1} \leq \alpha_{\ell_2} \leq \dots \leq \alpha_{\ell_s}$ .

**Definition 2.** Let  $X^* \subset \bar{D}$  be a manifold of dimension  $m$ , with  $0 \leq m < d$ . The coefficient distribution  $\alpha$  is called *type- $m$   $X^*$ -quasi-monotone on  $D$* , if for all  $\ell = 1, \dots, n$  there exists an index  $k$  such that  $X^* \subset \bar{Y}_k$  and such that there is a type- $m$  quasi-monotone path  $P_{\ell, k}$  from  $Y_\ell$  to  $Y_k$ .

**Definition 3.** Let  $\Gamma \subset \partial D$ . The coefficient distribution  $\alpha$  is called *type- $m$   $\Gamma$ -quasi-monotone on  $D$* , if for all  $\ell = 1, \dots, n$  there exists a manifold  $X_\ell^* \subset \Gamma$  of dimension  $m$  and an index  $k$  such that  $X_\ell^* \subset \partial Y_k$  and such that there is a type- $m$  quasi-monotone path  $P_{\ell, k}$  from  $Y_\ell$  to  $Y_k$ .

Note that the above definitions generalize the notion of quasi-monotone coefficients introduced in [3]. Definition 2 will be used to formulate weighted (discrete) Poincaré type inequalities, whereas Definition 3 will be used in weighted (discrete)



**Fig. 1** Examples of quasi-monotone coefficients. The numbering of the regions is according to the relative size of the coefficients on these regions with the smallest coefficient in region  $Y_1$ . Note that the first case is quasi-monotone in the sense of [3], but the other three are not. The first three examples are type-1. The last example is type-0. The manifold  $X^*$  is shown in each case, together with a typical path in some of the cases.

Friedrichs type inequalities. In the latter case,  $\Gamma$  plays the role of the Dirichlet boundary, where the function under consideration is assumed to vanish. In Figure 1 we give some examples of coefficient distributions that satisfy Definition 2.

To formulate our results we define for any  $u \in H^1(D)$  the average

$$\bar{u}^{X^*} := \begin{cases} u(X^*) & \text{if } \dim(X^*) = 0, \\ \frac{1}{|X^*|} \int_{X^*} u \, ds & \text{otherwise,} \end{cases}$$

as well as the weighted norm and seminorm

$$\|u\|_{L^2(D),\alpha} := \left( \int_D \alpha |u|^2 dx \right)^{1/2} \quad \text{and} \quad |u|_{H^1(D),\alpha} := \left( \int_D \alpha |\nabla u|^2 dx \right)^{1/2}.$$

**Lemma 1 (weighted Poincaré inequality).** *Let the coefficient  $\alpha$  be type- $(d-1)$   $X^*$ -quasi-monotone on  $D$  with the  $(d-1)$ -dimensional manifold  $X^*$ . For each index  $\ell = 1, \dots, n$ , let  $P_{\ell,k}$  be the path in Definition 2 with  $X^* \subset \bar{Y}_k$ , and let  $C_{\ell,k}^P > 0$  be the best constant in the inequality*

$$\|u - \bar{u}^{X^*}\|_{L^2(Y_\ell)}^2 \leq C_{\ell,k}^P \text{diam}(D)^2 |u|_{H^1(P_{\ell,k})}^2 \quad \text{for all } u \in H^1(P_{\ell,k}). \quad (2)$$

*Then there exists a constant  $C^P \leq \sum_{\ell=1}^n C_{\ell,k}^P$  independent of  $\alpha$  and  $\text{diam}(D)$  such that*

$$\|u - \bar{u}^{X^*}\|_{L^2(D),\alpha}^2 \leq C^P \text{diam}(D)^2 |u|_{H^1(D),\alpha}^2 \quad \text{for all } u \in H^1(D).$$

*Proof.* Let us fix one of the subregions  $Y_\ell$  and suppose without loss of generality that  $\int_{X^*} u \, ds = 0$  and that  $\text{diam}(D) = 1$ . Due to the assumption on  $\alpha$ , we have  $\|u\|_{L^2(Y_\ell),\alpha}^2 = \alpha_\ell \|u\|_{L^2(Y_\ell)}^2$ . Combining this identity with inequality (2) and using that the coefficients are monotonically increasing in the path from  $Y_\ell$  to  $Y_k$ , we obtain

$$\|u\|_{L^2(Y_\ell),\alpha}^2 \leq C_{\ell,k}^P \alpha_\ell |u|_{H^1(P_{\ell,k})}^2 \leq C_{\ell,k}^P |u|_{H^1(P_{\ell,k},\alpha)}^2 \leq C_{\ell,k}^P |u|_{H^1(D),\alpha}^2.$$

The proof is completed by adding up the above estimates for  $\ell = 1, \dots, n$ .

*Remark 1.* Obviously, inequality (2) follows from the standard Poincaré type inequality  $\|u - \bar{u}^{X^*}\|_{L^2(P_{\ell,k})}^2 \leq C |u|_{H^1(P_{\ell,k})}^2$  for all  $u \in H^1(P_{\ell,k})$ , with some constant  $C$  depending on  $P_{\ell,k}$  and on  $X^*$ , cf. [16, Sect. A.4]. However, this may lead to a sub-optimal constant. In general, the constants  $C_{\ell,k}^P$  depend on the choice of the manifold  $X^*$ , as well as on the number, shape, and size of the subregions  $Y_\ell$ . In Section 3, we give a bound of  $C_{\ell,k}^P$  in terms of local Poincaré constants on the individual subregions  $Y_\ell$  to make this dependency more explicit.

On the other hand, if  $X^*$  is a manifold of dimension less than  $d - 1$  (i.e. an edge or a point), the inequality in Lemma 1 does not hold for all functions  $u \in H^1(D)$ . However, there is a discrete analogue for finite element functions which holds under some geometric assumptions on the subregions  $Y_\ell$ , cf. [16, Sect. 4.6], in particular Lemma 4.15, Lemma 4.21, and the discussion thereafter.

Let  $\{\mathcal{T}_h(D)\}$  be a family of quasi-uniform, simplicial triangulations of  $D$  that are geometrically conforming with mesh width  $h$ . By  $V^h(D)$  we denote the space of continuous piecewise linear functions with respect to the elements of  $\mathcal{T}_h(D)$ . Note that we do not prescribe any boundary conditions. We further assume that the fine mesh  $\mathcal{T}_h(D)$  resolves the interfaces between the subregions  $Y_\ell$ .

**Assumption A.1** (cf. [16, Assumption 4.3]). There exists a parameter  $\eta$  with  $h \leq \eta \leq \text{diam}(D)$  such that each subregion  $Y_\ell$  is the union of a few simplices of diameter  $\eta$ , and the resulting coarse mesh is globally conforming on all of  $D$ .

Before stating the next lemma, we define the function

$$\sigma^\delta(x) := \begin{cases} (1 + \log(x)) & \text{for } \delta = 2, \\ x & \text{for } \delta = 3. \end{cases} \quad (3)$$

**Lemma 2 (weighted discrete Poincaré inequality).** *Let Assumption A.1 hold and let  $\alpha$  be type- $m$   $X^*$ -quasi-monotone on  $D$  with the manifold  $X^*$  having dimension  $m < d - 1$ . If  $m = 1$ , assume furthermore that  $X^*$  is an edge of the coarse triangulation in Assumption A.1. For each  $\ell = 1, \dots, n$ , let  $P_{\ell,k}$  be the path in Definition 2 with  $X^* \subset \bar{Y}_k$  and let  $C_{\ell,k}^{P,m} > 0$  be the best constant independent of  $h$  such that*

$$\|u - \bar{u}^{X^*}\|_{L^2(Y_\ell)}^2 \leq C_{\ell,k}^{P,m} \sigma^{d-m}\left(\frac{\eta}{h}\right) \text{diam}(D)^2 |u|_{H^1(P_{\ell,k})}^2 \quad \text{for all } u \in V^h(P_{\ell,k}). \quad (4)$$

*Then, there exists a constant  $C^{P,m} \leq \sum_{\ell=1}^n C_{\ell,k}^{P,m}$ , independent of  $h$ , of  $\alpha$ , and of  $\text{diam}(D)$  such that*

$$\|u - \bar{u}^{X^*}\|_{L^2(D),\alpha}^2 \leq C^{P,m} \sigma^{d-m}\left(\frac{\eta}{h}\right) \text{diam}(D)^2 |u|_{H^1(D),\alpha}^2 \quad \text{for all } u \in V^h(D).$$

*Proof.* The proof follows the same arguments as that of Lemma 1, but using (4) instead of inequality (2).

We remark that the existence of the constants  $C_{\ell,k}^{P,m}$  fulfilling inequality (4) will follow from the results summarized in [16, Sect. 4.6] and from our investigation in Section 3. For simplicity, let us also define  $\sigma^1 \equiv 1$  and  $C^{P,d-1} := C^P$ .

Similar inequalities than those in Lemmas 1 and 2 can also be proved, if  $u$  vanishes on part of the boundary of  $D$ .

**Lemma 3 (weighted (discrete) Friedrichs inequality).** *Let  $\Gamma \subset \partial D$  and let  $\alpha$  be type- $m$   $\Gamma$ -quasi-monotone on  $D$  (according to Definition 3). If  $m = d - 1$ , there exists a constant  $C^F = C^{F,d-1}$  independent of  $\alpha$  and of  $\text{diam}(D)$  such that*

$$\|u\|_{L^2(D),\alpha}^2 \leq C^F \text{diam}(D)^2 |u|_{H^1(D),\alpha}^2 \quad \text{for all } u \in H^1(D) \text{ with } u|_{\Gamma} = 0.$$

If  $m < d - 1$  and Assumption A.1 holds such that each  $X_\ell^*$  in Definition 3 is either a vertex or an edge of the coarse mesh in Assumption A.1, then

$$\|u\|_{L^2(D),\alpha}^2 \leq C^{F,m} \sigma^{d-m} \left(\frac{\eta}{h}\right) \text{diam}(D)^2 |u|_{H^1(D),\alpha}^2 \quad \text{for all } u \in V^h(D) \text{ with } u|_{\Gamma} = 0,$$

with  $C^{F,m}$  independent of  $h$ , of  $\alpha$ , and of  $\text{diam}(D)$ .

Similarly to the previous lemmas, the constants  $C^{F,m}$  in Lemma 3 are bounded by the sum of some constants  $C_{\ell,k}^{F,m}$  in standard Friedrichs type inequalities on the paths  $P_{\ell,k}$  in Definition 3. However, we will not discuss this further.

### 3 Explicit dependence on geometrical parameters

In this section we will study the dependence of the constants  $C_{\ell,k}^{P,m}$  (and consequently  $C^{P,m}$ ) in the above lemmas on the choice of  $X^*$  and on the number, size and shape of the regions  $Y_\ell$  (in particular the ratio  $\text{diam}(D)/\eta$ ). In [11, §3] the dependence on the geometry of the subregions is made more explicit. The lemmas presented there are in fact special cases of Lemmas 1 and 2 here.

First, we show that bounds for the constants  $C_{\ell,k}^{P,m}$  can be obtained from inequalities on the individual subregions  $Y_\ell$ . Secondly, we will look at some examples.

**Lemma 4.** *Let  $\alpha$  be type- $m$   $X^*$ -quasi-monotone on  $D$  with  $0 \leq m \leq d - 1$ , and let  $P_{\ell_1, \ell_s}$  be any of the paths in Definition 2. If  $m < d - 1$ , let Assumption A.1 hold. If  $m = 1$  and  $d = 3$ , assume additionally that  $X^*$  is an edge of the coarse triangulation. For each  $i = 1, \dots, s$ , let  $C_{\ell_i}^{P,m}$  be the best constant, such that*

$$\|u - \bar{u}^X\|_{L^2(Y_{\ell_i})}^2 \leq C_{\ell_i}^{P,m} \sigma^{d-m} \left(\frac{\eta}{h}\right) \text{diam}(Y_{\ell_i})^2 |u|_{H^1(Y_{\ell_i})}^2 \quad \text{for all } u \in V^h(Y_{\ell_i}), \quad (5)$$

where  $X \subset \bar{Y}_{\ell_i}$  is any of the manifolds  $X_{i-1}$ ,  $X_i$  or  $X^*$  in Definition 2 (as appropriate), cf. [16, Sect. 4.6]. Then

$$C_{\ell_1, \ell_s}^{P,m} \leq 4 \left\{ \sum_{i=1}^s \frac{\text{meas}(Y_{\ell_i})}{\text{meas}(Y_{\ell_i})} \frac{\text{diam}(Y_{\ell_i})^2}{\text{diam}(D)^2} C_{\ell_i}^{P,m} \right\}.$$

If  $m = d - 1$  we can extend the result to the whole of  $H^1$ .

*Proof.* We give the proof for the case  $m = d - 1$ . The other cases are analogous. For convenience let  $X_s := X^*$ . Then, telescoping yields

$$\|u - \bar{u}^{X^*}\|_{L^2(Y_{\ell_1})} \leq \|u - \bar{u}^{X_1}\|_{L^2(Y_{\ell_1})} + \sum_{i=2}^s \sqrt{\text{meas}(Y_{\ell_1})} |\bar{u}^{X_{i-1}} - \bar{u}^{X_i}|.$$

Due to (5), the first term on the right hand side is bounded by  $\sqrt{C_{\ell_1}^{P,m}} \text{diam}(Y_{\ell_1}) |u|_{H^1(Y_{\ell_1})}$ . For a fixed  $i$ , we can also conclude from inequality (5) that

$$\begin{aligned} |\bar{u}^{X_{i-1}} - \bar{u}^{X_i}|^2 &\leq \frac{2}{\text{meas}(Y_{\ell_i})} \left( \|\bar{u}^{X_{i-1}} - u\|_{L^2(Y_{\ell_i})}^2 + \|u - \bar{u}^{X_i}\|_{L^2(Y_{\ell_i})}^2 \right) \\ &\leq \frac{4}{\text{meas}(Y_{\ell_i})} C_{\ell_i}^{P,m} \text{diam}(Y_{\ell_i})^2 |u|_{H^1(Y_{\ell_i})}^2. \end{aligned}$$

An application of Cauchy's inequality (in  $\mathbb{R}^s$ ) yields the final result.

Let us look at some examples now. Firstly, if Assumption A.1 holds with constant  $\eta \gtrsim \text{diam}(D)$  (e. g. in Figure 1(a)), then  $n = \mathcal{O}(1)$  and each path  $P_{\ell,k}$  in Definition 2 contains  $\mathcal{O}(1)$  subregions. If we choose  $X^*$  to be a vertex, edge or face of the coarse triangulation in Assumption A.1, then by standard arguments  $C_{\ell}^{P,m} = \mathcal{O}(1)$  in (5) for all  $\ell = 1, \dots, n$ . Hence, it follows from Lemma 4 that the constants  $C^{P,m}$  in Lemmas 1–2 are all  $\mathcal{O}(1)$ .


Before we look to more complicated examples, which involve in particular long, thin regions, let us first derive two auxiliary results.

1. The middle region  $Y_3$  in Figure 1(b) is long and thin if  $\eta \ll \text{diam}(Y_3)$ . With  $X^*$  as given in the figure, one can show that (5) holds with  $C_3^{P,1} = \mathcal{O}(1)$ , independent of  $\eta$  and  $\text{diam}(Y_3)$ . This is not so surprising as  $\text{diam}(X^*) \simeq \text{diam}(Y_3)$ .
2. The region  $Y_8$  in Figure 1(c) has essentially the same shape, but here  $X^*$  has diameter  $\eta \ll \text{diam}(Y_8)$ . Nevertheless, one can show that (5) holds with  $C_8^{P,1} = \mathcal{O}(1)$ , independent of  $\eta$  and  $\text{diam}(Y_8)$ . (This result can be obtained by sub-dividing  $Y_8$  into small quadrilaterals of sidelength  $\eta$  and applying Lemma 4).

In Figures 1 and 2, we denote by  $H$  the sidelength of  $D$  (thus,  $H \simeq \text{diam}(D)$ ). We view  $\eta$  (if displayed) as a varying parameter  $\leq H$ , with the other parameters fixed.

**Figure 1(b).** As we have just discussed,  $C_{3,3}^{P,1} = C_3^{P,1} = \mathcal{O}(1)$ . Similarly,  $C_{2,2}^{P,1} = C_2^{P,1} = \mathcal{O}(1)$ . To obtain  $C_{1,3}^{P,1} = \mathcal{O}(1)$  we use  $\|u - \bar{u}^{X^*}\|_{L^2(Y_1)}^2 \leq \|u - \bar{u}^{X^*}\|_{L^2(P_{1,3})}^2$  and apply a standard Poincaré inequality (rather than resorting to Lemma 4 which would yield a pessimistic bound). Hence, Lemma 1 holds with  $C^{P,1} = \mathcal{O}(1)$ .

**Figure 1(c).** Despite the fact that  $C_1^{P,1} = \mathcal{O}(1)$  and  $C_8^{P,1} = \mathcal{O}(1)$ , the constant  $C_{1,8}^{P,1}$  is not  $\mathcal{O}(1)$ : Since  $\text{diam}(Y_1) \sim H$ , Lemma 4 yields  $C_{1,8}^{P,1} \lesssim \frac{H^2}{H^2} \frac{H^2}{H^2} + \frac{H^2}{H\eta} \frac{H^2}{H^2} = \mathcal{O}\left(\frac{H}{\eta}\right)$ . We easily convince ourselves that this is the worst constant  $C_{\ell,k}^{P,1}$ , for all  $\ell = 1, \dots, 9$  (e. g.,  $C_{3,9}^{P,1} = \mathcal{O}(1)$ ), and so we obtain  $C^{P,1} = \mathcal{O}\left(\frac{H}{\eta}\right)$ .



**Fig. 2** More examples (with  $\alpha_1 \ll \alpha_2$ ): The first two examples are quasi-monotone of type-1 and type-0, respectively.  $X^*$  is shown in each case. The examples in (c) and (d) are not quasi-monotone.

**Figure 1(d).** This coefficient distribution is only type-0 quasi-monotone and so we cannot apply Lemma 1, but by applying Lemma 4 we find that  $C_{7,8}^{P,0} = \mathcal{O}(1)$  and all the other constants are no worse. So in contrast to Case (c), we can show that the constant  $C^{P,0}$  in Lemma 2 is  $\mathcal{O}(1)$  in this case. The crucial difference is not that  $\alpha$  is type-0 here, but that  $\text{diam}(Y_8) = \mathcal{O}(H)$  and  $\text{diam}(Y_9) = \mathcal{O}(H)$ .

The examples in Figure 2 are further, typical test cases used in the literature.

**Figure 2(a).** To obtain a sharp bound for  $C^{P,1}$ , it is better here to treat all the regions where  $\alpha = \alpha_1$  as one single region  $Y_1$ , slightly modifying the proof of Lemma 1. Then  $C_{1,2}^{P,1} = \mathcal{O}(1)$  (standard Poincaré on  $D$ ). Due to a tricky overlapping argument that can be found in the Appendix of [10],  $C_{2,2}^{P,1} = \mathcal{O}(1)$ . Thus,  $C^{P,1} = \mathcal{O}(1)$ . If there are  $p$  distinct values in the inclusions, the constant  $C^{P,1}$  depends (linearly) on  $p$ . This should be compared with one of the main results in [5], where a similar Poincaré inequality is proved with a constant depending on the number of inclusions.

**Figure 2(b).** For each region  $Y_\ell$  we have  $C_\ell^{P,0} = C_\square^{P,0} = \mathcal{O}(1)$ , cf. [16]. For a moment, let us restrict on the regions where the coefficient is  $\alpha_1$  and group them into  $T := \frac{H}{2\eta}$  concentric layers starting from the two centre squares touching  $X^*$  where  $\alpha = \alpha_1$ . Obviously, for  $t = 1, \dots, T$ , layer  $t$  contains  $2t - 2$  regions where  $\alpha = \alpha_1$ . Each region in layer  $t$  can be connected to one of the two centre squares by a type-0 quasi-monotone path of length  $t$ . By Lemma 4,  $C_{\ell,k}^{P,0} \leq 4 \sum_{j=1}^t \frac{\eta^2}{H^2} C_\square^{P,0} = 4t \frac{\eta^2}{H^2} C_\square^{P,0}$  for all the regions  $Y_\ell$  in layer  $t$  where  $\alpha = \alpha_1$ . The same bound holds for the regions where  $\alpha = \alpha_2$ . Summing up these bounds over all regions and all layers, we obtain

$$C^{P,0} \leq 2 \sum_{t=1}^T (2t - 2) 4t \frac{\eta^2}{H^2} C_\square^{P,0} = 16 \frac{\eta^2}{H^2} \frac{T^3 - T}{3} = \mathcal{O}\left(\frac{H}{\eta}\right).$$

Equivalently, as there are  $n_\times = \mathcal{O}\left(\frac{H}{\eta}\right)^2$  crosspoints in this example,  $C^{P,0} = \mathcal{O}(\sqrt{n_\times})$ .

**Figure 2(c).**  $\alpha$  is not quasi-monotone in this case, and indeed Lemmas 1–3 do not hold. For example, if we choose  $X^*$  as shown, then it suffices to choose  $u$  to be the continuous function that is equal to  $2(x_1 - \frac{1}{4})$  for  $\frac{1}{4} \leq x_1 \leq \frac{3}{4}$  and constant otherwise, to obtain a counter example in  $V^h(D) \subset H^1(D)$  that satisfies  $\bar{u}^{X^*} = 0$ . We have  $\|u\|_{L^2(D),\alpha}^2 = \frac{\alpha_1}{6} + \frac{\alpha_2}{4}$  and  $|u|_{H^1(D),\alpha}^2 = 2\alpha_1$ , and so the constant  $C^{P,1}$  in Lemma 1 blows up with the contrast  $\frac{\alpha_2}{\alpha_1}$ . It is impossible to find  $X^*$  such that Lemma 2 holds.

**Figure 2(d).** Again  $\alpha$  is not quasi-monotone and Lemmas 1–3 do not hold on all of  $D$ . However, by choosing suitable (energy-minimising) coarse space basis functions in two-level Schwarz methods (cf. [6, 14, 9]), it often suffices to be able to apply Lemmas 1–3 on  $D' := Y_1 \cup Y_2 \cup Y_3$ . Since  $\alpha$  is type-1 quasi-monotone on  $D'$ , e.g. Lemma 1 holds for  $u \in H^1(D')$  and it is easy to verify that  $C^{P,1} = \mathcal{O}(1)$ .

For some 3D examples see [11]. The constants  $C^{F,m}$  in Lemma 3 behave similarly. Finally, for applications of weighted Poincaré–Friedrichs inequalities, such as Lemmas 1–3, in the analysis of FETI and two-level Schwarz methods see [11, 15].

## References

1. J. H. Bramble and J. Xu. Some estimates for a weighted  $L^2$  projection. *Math. Comp.*, 56:463–476, 1991.
2. C. R. Dohrmann, A. Klawonn, and O. B. Widlund. Domain decomposition for less regular subdomains: Overlapping Schwarz in 2d. *SIAM J. Numer. Anal.*, 46:2153–2168, 2008.
3. M. Dryja, M. V. Sarkis, and O. B. Widlund. Multilevel Schwarz methods for elliptic problems with discontinuous coefficients in three dimensions. *Numer. Math.*, 72:313–348, 1996.
4. Y. Efendiev and J. Galvis. Domain decomposition preconditioners for multiscale problems. Preprint, Texas A&M University, 2009. submitted.
5. J. Galvis and Y. Efendiev. Domain decomposition preconditioners for multiscale flows in high contrast media. Preprint, Texas A&M University, 2009. submitted.
6. I. G. Graham, P. Lechner, and R. Scheichl. Domain decomposition for multiscale PDEs. *Numer. Math.*, 106:589–626, 2007.
7. A. Klawonn, O. Rheinbach, and O. B. Widlund. An analysis of a FETI-DP algorithm on irregular subdomains in the plane. *SIAM J. Numer. Anal.*, 46:2484–2504, 2008.
8. A. Klawonn and O. B. Widlund. FETI and Neumann-Neumann iterative substructuring methods: connections and new results. *Comm. Pure Appl. Math.*, 54:57–90, 2001.
9. J. Van lent, R. Scheichl, and I. G. Graham. Energy minimizing coarse spaces for two-level Schwarz methods for multiscale PDEs. *Numer. Lin. Alg. Appl.*, 16:775–799, 2009.
10. C. Pechstein and R. Scheichl. Analysis of FETI methods for multiscale PDEs. *Numer. Math.*, 111:293–333, 2008.
11. C. Pechstein and R. Scheichl. Analysis of FETI methods for multiscale PDEs - Part II: Interface variation. BICS Preprint 7/09, University of Bath, UK, 2009. submitted.
12. C. Pechstein and R. Scheichl. Scaling up through domain decomposition. BICS Preprint 1/09, University of Bath, UK, 2009. to appear in *Appl. Anal.*
13. M. Sarkis. Nonstandard coarse spaces and Schwarz methods for elliptic problems with discontinuous coefficients using non-conforming elements. *Numer. Math.*, 77:383–406, 1997.
14. R. Scheichl and E. Vainikko. Additive Schwarz and aggregation-based coarsening for elliptic problems with highly variable coefficients. *Computing*, 80:319–343, 2007.
15. R. Scheichl, P. S. Vassilevski, and L. Zikatanov. Two-level Schwarz with non-matching coarse grids. Preprint, Lawrence Livermore National Labs, 2009.
16. A. Toselli and O. Widlund. *Domain Decomposition Methods – Algorithms and Theory*. Springer, Berlin, 2005.
17. J. Xu and Y. Zhu. Uniform convergent multigrid methods for elliptic problems with strongly discontinuous coefficients. *Math. Models Methods Appl. Sci.*, 18:1–29, 2008.
18. Yunrong Zhu. Domain decomposition preconditioners for elliptic equations with jump coefficients. *Numer. Linear Algebra Appl.*, 15:271–289, 2008.

## Latest Reports in this series

### 2009

- |         |  |               |
|---------|--|---------------|
| 2009-01 | Clemens Pechstein and Robert Scheichl<br><i>Scaling Up through Domain Decomposition</i>  | January 2009  |
| 2009-02 | Clemens Hofreither, Ulrich Langer and Satyendra Tomar<br><i>Boundary Elements Simulation of Linear Water Waves in a Model Basin</i>                          | February 2009 |
| 2009-03 | Huidong Yang and Walter Zulehner<br><i>Numerical Simulation of Fluid-Structure Interaction Problems on Hybrid Meshes with AMG</i>                            | April 2009    |
| 2009-04 | Clemens Pechstein and Robert Scheichl<br><i>Analysis of FETI Methods for Multiscale PDEs - Part II: Interface Variation</i>                                  | April 2009    |
| 2009-05 | Alexsandr M. Matsokin and Sergey V. Nepomnyaschikh<br><i>Domain Decomposition Preconditioning for Well Models for Reservoir Problems</i>                     | June 2009     |
| 2009-06 | Helmut Gfrerer<br><i>Generalized Penalty Methods for a Class of Convex Optimization Problems with Pointwise Inequality Constraints</i>                       | July 2009     |
| 2009-07 | Ulrich Langer, Huidong Yang and Walter Zulehner<br><i>A Grid-enabled Solver for the Fluid-Structure Interaction (FSI) Problem</i>                            | August 2009   |
| 2009-08 | Stefan Takacs and Walter Zulehner<br><i>Multigrid Methods for Elliptic Optimal Control Problems with Neumann Boundary Control</i>                            | October 2009  |
| 2009-09 | Dylan M. Copeland and Ulrich Langer<br><i>Domain Decomposition Solvers for Nonlinear Multiharmonic Finite Element Equations</i>                              | November 2009 |
| 2009-10 | Huidong Yang and Walter Zulehner<br><i>A Newton Based Fluid-structure Interaction (FSI) Solver with Algebraic Multigrid Methods (AMG) on Hybrid Meshes</i>   | November 2009 |
| 2009-11 | Peter Gruber, Dorothee Knees, Sergiy Nesenenko and Marita Thomas<br><i>Analytical and Numerical Aspects of Time-dependent Models with Internal Variables</i> | November 2009 |
| 2009-12 | Clemens Pechstein and Robert Scheichl<br><i>Weighted Poincaré Inequalities and Applications in Domain Decomposition</i>                                      | November 2009 |

From 1998 to 2008 reports were published by SFB013. Please see

<http://www.sfb013.uni-linz.ac.at/index.php?id=reports>

From 2004 on reports were also published by RICAM. Please see

<http://www.ricam.oeaw.ac.at/publications/list/>

For a complete list of NuMa reports see

<http://www.numa.uni-linz.ac.at/Publications/List/>